

Community Drug Distributor (CDD) Job Aid

Integrated Community Case Management (ICCM)




**Athor ci tueny-tueny ka nguan mat nhim bi CDD miith kony baaic
Warak tou anguen ma tich CDD**




Ruun

How old is the child?



2 to 11 months

**Peei karou bi la peei thiarku tok
Duatde ariou-laate apaar wang aciilo**



1 to 5 years

**Ruon tok bi la run kadhiec
Run-aciilo laate run abiic**

What is the child's sex?



Boy

Dhok
Nyithok



Girl


Nya
Nyakou



Not able to wake

Aci gek piny


Ungen piny



Convulsions

Aci thook

Atalhogo



Chest in-drawing

Puou thok aci lony thin

Ngeete, teel kii ngo



Not able to drink or breastfeed

Acie dek piu Acie thuat

Bimatha –Bidhodho



Vomiting everything

Akang ngak wei e ben

Ungok kuan gini came cang




Neck stiffness

Aci cik yeth

Ngut he talhogo


Atoc panakim Ahor parakim

If any danger sign refer to PHCU/PHCC




Atuoch guop Leeth rook

Assess for fever



Does the child have a fever?

**Meth neng atuoc
Nyithiin reeh lieth**



**Leng niin kadah kek atuoc
Yiere niniin adih**

For how long?


Aweei apei Yii yong yong

Assess for fast breathing and cough



Does the child
have a cough?

Meth eh hol
Nyithin uwlo




Does the child have fast breathing?

Meth eh wei apei
Nyithin uyii kerio



Yac Thou yec

Assess for diarrhoea



Watery diarrhoea

Yac piu
Thou-yec pii




Bloody diarrhoea

Yac riem
Thou-yec remo



Skin pinch test

Aci guop thok piu
Pii athum eguobe



For how long?


Leng niin kdah kek yac
Yiere niniin adih

Aci door Adoor kerayo juet

Assess for SAM



MUAC tape



Pedal Oedema

Aci buot cook

Tiene-kuot

Tuany ril yic Tou me yee tek

Refer for severe illness




Atuoc niin kadhoro leh tueng
Leeth rok niin abiberio ucidhe
wang malo



Fever for 7 days or more



Yac piu niin thiar ku nguan leh tueny
Thou yech niin apar wang anguen
ucidhe wang malo





Watery diarrhoea for
14 days or more




Bloody diarrhoea

Yac riem
Thou yac remo



Tuany door ril yic Tou me yee tek


Refer SAM



Gam wal panakim tene meth atuoc

Chup yeen parakim leeth rook


Treatment for fever



Gam wal panakim tene meth hol ku weei apei

Cup yeen parakim yier nyithin mu uulo yii yonge yong


Treatment for fast breathing and cough and chest in-drawing



AMOXICILLIN TABLETS FOR ORAL SUSPENSION USP 125 MG
2 X 10 TABLETS

Age: From 2 months to 11 months


Time	Day 1	Day 2	Day 3	Day 4	Day 5
	2 tabs	2 tabs	2 tabs	2 tabs	2 tabs
	2 tabs	2 tabs	2 tabs	2 tabs	2 tabs



AMOXICILLIN TABLETS FOR ORAL SUSPENSION USP 125 MG
3 X 10 TABLETS

Age: From 12 months to 59 months (1-5 years)

Time	Day 1	Day 2	Day 3	Day 4	Day 5
	3 tabs	3 tabs	3 tabs	3 tabs	3 tabs
	3 tabs	3 tabs	3 tabs	3 tabs	3 tabs



Gam wal panakim tene meth yac


Cup yeen parakim thou yac

Treatment for diarrhoea



**Awai ci mat ke
thugar**

**Kat melou man mat
ke cukar**



Piu path

Pii-me kuel






1 2 3 4 5 6 7 8 9 10



1 2 3 4 5 6 7 8 9 10

Gam wal panakim tene meth ci door Cup yeen ariou yier nyithiin ma adoor

Pre-referral treatment for SAM



Do not give ORS
and zinc to
children who are
malnourished




Duk wal ka gem tene meth ci door
Keri chup yeeni yier nyithiin ma
adoor


Caregiver Health Messages






Sleep under a mosquito net





Breast feed children for 2 years or more
Feed your child only breast milk for the first 6 months of life





Take children for immunizations






Wash hands with soap and water





Drink safe water





Use latrines





Give children different nutritious food to eat

