




**Peei karou bi la peei thiarku tok  
Duatde ariou-laar apaar wang aciilo**

Child 2-11 months



**Ruon tok bi la run kadhiec**  
**Run-aciilo laat run abiic**

Child 1 to 5 years




**Aci thook**  
**Atalhogo**  
Convulsion



**Aci gek piny**  
**Ungen piny**

Abnormally sleepy / not able to wake



**Puou thok aci lony thin  
Ngeete, teel kii ngo**

Chest in-drawing





**Acie dek piu Acie thuat  
Bimatha –Bidhodho**

Not able to drink or breastfeed



**Akang ngak wei e ben  
Ungok kuan gini came cang**


Vomiting everything



**Aci cik yeth**  
**Ngut he talhogo**

Neck stiffness






**Atoc panakim  
Ahor parakim**


Danger sign refer to PHCU/C











Fast breathing and cough





Respiratory timer



Respiratory beads 2-11 months





Respiratory beads child 1 to 5 years





**Yac piu Yac riem**  
**Thou-yec pii Thou-yec remo**  
Watery diarrhoea and bloody diarrhoea





Skin pinch test showing dehydration






Sunken eyes from dehydration




Children with SAM






Assessment pitting oedema both feet



MUAC tape



1 2 3 4 5 6 7




**Atuoc niin kadhoro leh tueng  
Leeth rok niin abiberio ucidhe wang malo**

Fever 7 days or more





Respiratory timer and beads




**Yac piu niin thiar ku nguan leh tueny**  
**Thou yech niin apar wang anguen ucidhe wang malo**  
Watery diarrhoea for 14 days or more



**Yac riem**  
**Thou yac remo**  
Bloody diarrhoea





**OTHER DISEASE  
GO TO PHCU/C**



**Tuany ril yic  
Tou me yee tek**


Severe illness refer PHCU/C












1

2

3



1



2













3



# AMOXICILLIN TABLETS FOR ORAL SUSPENSION USP 125 MG

2 X 10 TABLETS

Age: From 2 months to 11 months					
Time	Day 1	Day 2	Day 3	Day 4	Day 5
	2 tabs	2 tabs	2 tabs	2 tabs	2 tabs
	2 tabs	2 tabs	2 tabs	2 tabs	2 tabs

1

2



3


4

5

# AMOXICILLIN TABLETS FOR ORAL SUSPENSION USP 125 MG

3 X 10 TABLETS

Age: From 12 months to 59 months (1-5 years)					
Time	Day 1	Day 2	Day 3	Day 4	Day 5
	3 tabs	3 tabs	3 tabs	3 tabs	3 tabs
	3 tabs	3 tabs	3 tabs	3 tabs	3 tabs



1


2

3


4

5












Infant 2 to 6 months Zinc ½ tablet for 10 days










NEVER give Zinc or ORS to children with SAM









Breastfeed for 2 years








Wash hands





Boil water for 1 minute—drink safe water





Use pit latrine



# Instructional Pictures for ICCM Training of Community Drug Distributors





1. Infant 2-11 months
2. Child 1-5 years
3. Convulsion
4. Abnormally sleepy or not able to wake
5. Chest in-drawing
6. Not able to breastfeed or drink
7. Vomiting everything
8. Neck stiffness
9. Red Danger Sign Triangle with PHCU/C site
10. Photograph of PHCC
11. Fever
12. CDD touching forehead of child with fever
13. Fast breathing and cough
14. Respiratory timer
15. Respiratory beads for infants 2-11 months
16. Respiratory beads for children 1-5 years.
17. Watery diarrhoea and diarrhoea with blood in stool
18. Skin pinch test of dehydrated child
19. Sunken eyes of dehydrated child
20. Children with SAM
21. Pedal oedema
22. Measuring with MUAC tape
23. Fever with 7 suns
24. Respiratory timer and both respiratory beads with red beads circled
25. Watery diarrhoea with 14 suns
26. Bloody diarrhoea
27. Severe illness Yellow Triangle with PHCU/C site
28. Infant with SAM measured with MUAC tape
29. Red MUAC tape reading
30. Malnutrition referral triangle and OTP site
31. AS+AQ PINK box and 1 tablet 3 days
32. AS+AQ PURPLE box and 1 tablet 3 days
33. Amoxicillin PINK box 2 tablets twice a day for 5 days
34. Amoxicillin GREEN box 3 tablets twice a day for 5 days
35. ORS solution 3 packets
36. 1 litre of water container
37. Zinc tablets 1/2 tablet for 10 days
38. Zinc tablets 1 tablet for 10 days
39. SAM triangles and AS+AQ and amoxicillin
40. NEVER give ORS or Zinc to children with SAM
41. Give first dose of medicine with caregiver
42. Sleep under bed net
43. Breastfeed for 2 years
44. Eat a variety of nutritious foods
45. Wash hands with soap and water or ash
46. Drink safe water
47. Use latrine
48. Immunize children
49. Cover