

Understanding Child Health in the Context of COVID-19: Experiences from Bangladesh, Pakistan, & Zambia

April 23, 2020


COVID-19 Pandemic

David Hamer, MD

Dept. of Global Health, BU School of Public Health

Section of Infectious Diseases, Department of Medicine, BU School of Medicine

National Emerging Infectious Diseases Laboratory



Origin of the nCoV-2019?



Many species of bats harbor coronaviruses


 2017: a coronavirus 96% identical to the Wuhan coronavirus was identified in a horseshoe bat in Yunnan cave



Did pangolins spread the China coronavirus to people?

Genetic sequences of viruses isolated from the scaly animals are 99% similar to that of the circulating virus – but the work is yet to be formally published.

David Cyranoski



Early
transmission
dynamics in
Wuhan, China
of novel
coronavirusinfected
pneumonia.
NEJM 2020

Qun Li et al.

Figure 1. Onset of Illness among the First 425 Confirmed Cases of Novel Coronavirus (2019-nCoV)—Infected Pneumonia (NCIP) in Wuhan, China.

Figure 1. Epidemic Curve of the Confirmed Cases of Coronavirus Disease 2019 (COVID-19) 4000-Confirmed cases only 3500-By date of onset (n=44672) By date of diagnosis (n = 44672) 3000 of cases 1500 10 1000-500-2020 30,2020 2019 2019 2019 2019 2020 Jan 8, 2020 10, 2020 2000 200 B, 2020 M, 2020 5,2020 2020 8 8 4,2020 Jan 9, 2020 4 Unusual cases of pneumonia COVID-19 (3 in the same family) noticed by Huanan Seafood 2019-nCoV a Class B Jixian Zhang, MD, in HICWM Hospital. Market closed Identified notifiable disease WHO declares "public Dr Zhang reported unusual Wuhan Health Commission 2019-nCoV health emergency of pneumonia cases to the alerts National Health sequences Wuhan City International concern* local CDC Commission and China CDC. filest shut down shared and WHO is notified. 3 More cases of Another 15 pneumonia found in HICWM cities shut Active case finding 2019-nCoV test Lunar New Year Nationwide mandatory Hospital (for a total of 7) begins in Wuhan City kits first available down national holiday extended holiday

Wu & McGoogan JAMA 2020

WHO Situational Report April 22, 2020

- WHO risk assessment: global - very high
- US>Spain>Italy>China> Iran
- Globally 2,471,136 confirmed cases (73,920 new)
 - 169,006 deaths (6058 new)

Figure 4. Epidemic curve of confirmed COVID-19, by date of report and WHO region through 22 April 2020






Figure 1. Countries, territories or areas with reported confirmed cases of COVID-19, 22 April 2020



Global Trajectory of COVID-19 in Africa and India vs. High-Income Countries

COVID-19 in Africa United Nations

Figure 1.1 Days since 100th case: Africa's infection trajectory vs comparators




Source: : Based on data from Johns Hopkins University and Africa CDC, 12 April 2020

Enhancing capacity for COVID-19 diagnostics in Bangladesh

Child Health Research Foundation, Bangladesh
Prof. Samir K Saha and Dr. Senjuti Saha
23 April 2020
Child Health Task Force Webinar

COVID-19 in Bangladesh


3,772 cases, 120 deaths



Data from ourworldindata.org

Age distribution of COVID-19 cases in Bangladesh

The figure below is showing age distribution of reported confirmed COVID-19 cases (n=1,902), including deaths (n=58), Bangladesh.



Gaps in knowledge

- Like many other places, there is a general assumption that children are "resistant" to COVID-19
- No data on
 - symptoms
 - duration of illness
 - hospitalization
 - contact history
 - risk factors (comorbidities, nutritional status)
- No data on asymptomatic carriage

Child Health Research Foundation in Bangladesh

The objectives of our semi-national surveillance are:

- to describe epidemiology and estimate burden of vaccinepreventable disease
 - Meningitis, sepsis, pneumonia, diarrhea
- to establish a platform to measure vaccine needs and impact
- to characterize circulating bacterial and viral types AMR, serotypes, etc






CHRF's journey during the COVID-19 era – in progress


- March 24 Gov. of Bangladesh requested that we prepare to serve as a COVID-19 diagnostic lab
- March 24 29 We converted our primary research lab into COVID-19 diagnostic lab
- March 29 Official letter from GoB to become a COVID-19 lab
 - COVID-19 testing starts
- April 02 6 phone lines were set up for suspect COVID-19 patients and report delivery/guidance
- Currently
 - Test ~250/day (7% of Bangladesh COVID-19 data)
 - Provide onsite and offsite trainings to hospital on biosafety (PPE use), sample collection and transportation
 - Provide collection logistics (tubes, media/RNA shield/flocked swab) for swab collection

Overview of samples tested to date





Age distribution of CHRF COVID-19 +/- cases




Future plans (funded by the Bill and Melinda Gates Foundation)

Objective: To collect data on hospitalization, severity and long-term outcome of pediatric COVID-19 patients admitted in the two largest children's hospitals in Bangladesh

- Immediate-future plans
 - From 24 April, sample collection forms for symptoms and duration will be filled in majority of the collection sites
 - For children admitted in our hospital, we will conduct daily follow-up
 - From 26 April, we are starting a phone follow-up for pediatric cases prospectively
 - Data on comorbidity, duration of illness, contact history, hospitalization
 - Follow-up calls will continue to gather data on outcome up to 90 days following test results

We shall overcome



COVID-19 in Pakistan

Pediatric Perspective

Global Child Health Task Force

Dr. Asad Ali

Professor, Pediatric infectious Diseases

&

Dr. Faisal Mahmood Section Head, Adult Infectious Diseases

> AGA KHAN UNIVERSITY KARACHI, PAKISTAN

_gB **290**

Jan 1, 2020 - Apr 23, 2020

Confirmed Cases

10,513



Active Cases



Deaths

2.1%





Recoveries


22.2%


Last updated on


	Confirmed Cases	Active Cases	Deaths	Recoveries
AJK	51	28	0	23
Balochistan	552	376	8	168
GB	290	86	3	201
Islamabad	204	175	3	26
KPK	1,453	956	83	414
Punjab	4,590	3,742	58	790
Sindh	3,373	2,589	69	715

















Challenges/Opportunities

Limited infrasatructure if the disease turns more virulent in children

Community pediatricians remain at high risk – Need revised Standard
 Operating Procedures in busy outpatient clinics

• Children as a portal of community transmission, even if the disease is relatively less severe

Inclsuion of children in the vaccine trials.

Rise of COVID-19 in Africa





Africa CDC Dashboard

Cases **25,131**

Deaths **1,205**


Recoveries 6,576






Region All

Figure 1.3 Proportion of urban population living in slums, percent




Source: Based on data from UN Habitat, 2016

Figure 1.4 Access to household handwashing facilities



Source: Based on WASH data from WHO/UNICEF, 2017

Figure 1.8 HIV/AIDS, prevalence



The boundaries and names shown and the designations used on this map do not imply official endorsement or acceptance by the United Nations. Final boundary between


the Republic of Sudan and the Republic of South Sudan has not yet been determined

Source: Based on data from Global Burden of Disease Study, 2016

Figure 1.6 Chronic respiratory diseases, prevalence



Figure 1.9 With 1.8 average hospital beds per 1,000 people, hospital beds capacity across Africa is weak



Source: Index Mundi from World Health Organization, 2020

Table 1.2 Cost of COVID-19 medical supply response (gap) across Africa, by pandemic scenario

	A - Unmitigated	B - Mitigation using moderate physical distancing	C - Suppression using intense physical distancing	D - Suppression using intense and early physical distancing
Cost of COVID-19 medical supply response (gap) across Africa, 2020 (\$ billion)	\$446	\$335	\$189	\$44
Africa's current health expenditure allocation, 2020 (\$ billion)	\$138.87	\$138.87	\$138.87	\$138.87
Projected percentage increase in required health spending (%)	321.16	240.89	136.09	31.83

A: Unmitigated (worst case) - no intervention

B: Mitigation using moderate social distancing - Optimal outcome when epidemic is mitigated through interventions to limit contacts in general population including social distancing (45% reduction in contact rate)


C: Suppression using intense social distancing (1.6) - Introduce intense social distancing measures that reduce the contact rate in the general population by 75% once 1.6 deaths per million per week trigger is reached

D: Suppression using intense social distancing (0.2) - Introduce intense social distancing measures that reduce the contact rate in the general population by 75% once 0.2 deaths per million per week trigger is reached

Source: ECA cost estimates using Imperial College demand figures and various sources for prices (see annex I below)

Potential Impact on GDP in Africa

Figure 2.1 Expected drop in growth from COVID-19 impact, ECA estimates



Source: ECA estimates, 2020